



Role of fish in mixing and spreading nutrients in coastal waters revealed

April 7, 2022 | Steve Bates, University of Southampton

Topics: Ecosystems

A new study has shown how fish influence oceans' ecosystems in coastal regions, revealing for the first time the role they play in distributing heat, nutrients and oxygen that keep the system functioning.

Oceans are made up of multiple layers, ranging from lighter, warmer waters at the top to denser, cooler waters at the bottom. Ocean mixing is vital to move heat, oxygen, nutrients and pollutants between different layers and therefore plays a major role in how ecosystems can sustain life. Although it is well established that winds and tides supply the bulk of the energy that drives mixing, the contribution made by swimming organisms has not been understood until now.

In this new study, the University of Southampton, together with a team including the aquatic research institute Eawag, spent fifteen days monitoring water turbulence in the Ría de Pontevedra, a bay in the north-west coast of the Iberian Peninsula. The researchers used an instrument called microstructure profiler, which measures fluctuations of the ocean current velocities and temperature at very small distances.



Observing turbulence data.
(Photo: Remedios project)

The results, published in the journal *Nature Geoscience*, showed that elevated levels of turbulence and mixing occurred every night, comparable to turbulence caused by a major storm, despite the weather staying calm throughout the study. Using acoustic information from the ship-echosounder and samples collected with small fishing nets the research team were able to attribute the signal to the presence of fish aggregations, which gathered in the area at night. In particular, the small nets were full of recently spawned eggs of European anchovy, *Engraulis encrasicolus*, providing a strong evidence that the anchovies' frantic behaviour during spawning created the turbulence. .

“We believe that biological mixing was intense in our observations because the bay is highly stratified - temperature and other properties vary significantly at different depths,” explained Dr Bieito Fernández Castro, a research fellow at the University of Southampton who led the study. “Previous studies have suggested that biological turbulence causes little mixing because the circular motions of water that the fish generate while swimming are too small. This is certainly true in the open ocean, where temperature changes occur over tens of meters. However, we have shown that closer to land, where the layers change over a much shorter distance, the anchovies are able to mix them together,” he continued.

The study has revealed that whilst biological mixing may not be very important in the open ocean, it can be significant in coastal ecosystems, where a thriving marine life coexists with rapid vertical changes in the make-up of the ocean. The vertical mixing created by fish schools could impact the redistribution of temperature, nutrients and other important water constituents, like oxygen, which play a fundamental role in the functioning of the ecosystem upon which the fish themselves rely. The findings therefore highlight the capacity of living organisms to influence and reshape the physical environment where they live.

“The observation of how our anchovies drove mixing was totally fortuitous. We were set to study how turbulence affects marine life and we end up showing, for the first time, that marine life can influence ocean turbulence, which in turn influences marine life!” Dr Fernández Castro concluded.

The study formed part of the [REMEDIOS research project](#), which is led by the University of Vigo (Spain) with partners including the University of Southampton and investigates the role

of mixing in phytoplankton growth.

Biomixing

Credits: Remedios Proyecto, Espe Broullón

Cover picture: Research Vessel Ramón Margalef (Spanish Institute of Oceanography) departing from Vigo harbour. (Photo: Remedios project)

Original publication

Bieito Fernández Castro, Marian Peña, Enrique Nogueira, Miguel Gilcoto, Esperanza Broullón, Antonio Comesaña, Damien Bouffard, Alberto C. Naveira Garabato, Beatriz Mouriño-Carballido (2022) Intense upper ocean mixing due to large aggregations of spawning fish, [Nature Geoscience](#)

Cooperations

University of Southampton, UK Instituto de Investigacions Marinas, Spain Instituto Espanol de Oceanografía, Spain Universidade de Vigo, Spain Eawag, Switzerland

Related Links

Research Project Remedios

Media information of the University of Southampton

Steve Bates
Media Relations Officer
University of Southampton
s.d.bates@soton.ac.uk
+44 7342 060429

Contact



Damien Bouffard

Tel. +41 58 765 2273

damien.bouffard@eawag.ch



Bärbel Zierl

Science editor

Tel. +41 58 765 6840

baerbel.zierl@eawag.ch

<https://www.eawag.ch/en/info/portal/news/news-archive/archive-detail/role-of-fish-in-mixing-and-spreading-nutrients-in-coastal-waters-revealed>