

Conversion product of painkiller more toxic than expected

April 28, 2020 | Stephanie Schnydrig Topics: Wastewater | Pollutants

The active ingredient diclofenac contained in various painkillers is hardly degraded at all in wastewater treatment plants and therefore accumulates in surface waters. There, amphipods convert diclofenac into a more toxic substance, namely diclofenac methyl ester, as shown in a study by the Swiss Federal Institute of Aquatic Science and Technology (Eawag).

Every day, several tons of pharmaceutical ingredients are released into bodies of water worldwide. These originate mainly from domestic wastewater, because most substances are excreted again after ingestion – in part unchanged. Since many wastewater treatment plants do not retain all substances without leaving any residues, the rest ends up in the aquatic environment. Even if the amounts are extremely small, they can adversely affect organisms. Previous studies have shown that the analgesic diclofenac has a harmful effect on the liver, kidneys and gills of fish.

Researchers led by Juliane Hollender, head of Eawag's Environmental Chemistry Department, are now presenting their findings: a conversion product of diclofenac, namely diclofenac methyl ester, may be even more toxic than the starting material itself. The research team recently published this finding in the journal "Environmental Science & Technology".

Amphipods convert active ingredient

Environmental chemists came to this conclusion with the help of two common types of amphipods. In the laboratory, post-doctoral researcher Qiuguo Fu exposed these to different concentrations of diclofenac for 24 hours. Admittedly, the concentrations chosen were several times higher than those found in the environment. Then she investigated how the substance behaves in the creatures' bodies.

T +41 58 765 55 11 F +41 58 765 50 28 info@eawag.ch www.eawag.ch



According to this study, an enzymatic reaction in the small creatures leads to the formation of the substance diclofenac methyl ester, which is acutely much more toxic than diclofenac.



The common amphipod, Gammarus pulex, converts diclofenac into the more acutely toxic diclofenac methyl ester. (Photo: Eawag)

Moreover, diclofenac methyl ester is poorly soluble in water and can therefore be excreted less well, meaning it can accumulate to a greater extent in the body. "I therefore consider the substance to be more dangerous than diclofenac," says Juliane Hollender. According to the environmental chemist, this type of chemical transformation occurred unexpectedly and should be considered in toxicological risk assessments for other substances. For according to initial studies, this biotransformation also occurs in higher organisms such as fish and humans.

The good news is that around a hundred wastewater treatment plants in Switzerland are currently being equipped with a fourth treatment stage to effectively remove micropollutants. "After this additional elimination step, diclofenac no longer appears in the water in increased concentrations," says Hollender.

Cover picture: pxhere

Original Publication

Fu, Q.; Fedrizzi, D.; Kosfeld, V.; Schlechtriem, C.; Ganz, V.; Derrer, S.; Rentsch, D.; Hollender, J. (2020) Biotransformation changes bioaccumulation and toxicity of diclofenac in aquatic organisms, *Environmental Science and Technology*, 54, 4400-4408, doi: 10.1021/acs.est.9b07127, Institutional Repository

Funding

Schweizer Nationalfonds, European Chemical Industry Council

info@eawag.ch www.eawag.ch



Contact



Juliane Hollender

Senior scientist / Group leader Tel. +41 58 765 5493 juliane.hollender@eawag.ch

https://www.eawag.ch/en/info/portal/news/news-archive/archive-detail/conversion-product-of-painkiller-more-toxic-than-expected

